

Notes on domain theory and topology

Frédéric Blanqui

17 August 2012

These notes have been written to complement the study of the very nice book of Gilles Dowek on “Proofs and Algorithms - An Introduction to Logic and Computability” [2] by Ying Jiang’s group of students at the Institute of Software of the Chinese Academy of Sciences (ISCAS) that met more or less every Tuesday from May to August 2012.

It introduces the notion of directed-complete partial order (dCPO) that is the basis of domain theory [1] and topology.

1 Directed-complete partial orders

Let \leq be an ordering relation on a set E .

The first chapter of Dowek’s book starts by introducing the following notions:

Definition 1 (Limit, complete ordering, continuity) An element $l \in E$ is the *limit* of an *increasing* sequence $u_0 \leq u_1 \leq \dots$ if l is the *least upper bound* (lub) of the set $\{u_0, u_1, \dots\}$.

The ordering \leq is *weakly complete* if every increasing sequence has a limit. It is *strongly complete* if every subset of E has a least upper bound.

A function $f : E \rightarrow E$ is *increasing* if for all $x, y \in E$, $x \leq y \Rightarrow f(x) \leq f(y)$.

In a weakly complete ordering \leq , a function $f : E \rightarrow E$ is *continuous* if it is increasing and if, for every increasing sequence $(u_i)_{i \geq 0}$, $\text{lub}\{f(u_i)\}_{i \geq 0} = f(\text{lub}\{u_i\}_{i \geq 0})$.

A set E equipped with an ordering relation \leq is also called a *partially ordered set* or *poset*. In a *totally ordered set*, every two elements are comparable.

An increasing function is more usually called *monotone*, *monotonic* or *isotone* (it is a morphism in the category of posets).

An increasing sequence u_0, u_1, \dots is also called an ω -*chain* (ω is the first infinite ordinal), a *chain* being a non-empty totally ordered set. Because of this restriction to ω -chains, Dowek’s notions of weak completeness and continuity are generally called ω -completeness (ω CPO) and ω -continuity.

Moreover, a strongly complete ordering is a *complete lattice*:

Definition 2 (Lattice) A *lattice* is a poset (E, \leq) such that every two elements have a least upper bound (lub) and a greatest lower bound (glb). A lattice is *complete* if every subset of E has a lub and a glb.

An easy induction shows that, in a lattice, every *non-empty finite* subset of E has a lub and a glb. It is complete if every empty and infinite subset has a lub and a glb.

Lemma 3 A strongly complete ordering is a complete lattice.

Proof. The fact that every subset has a lub follows from the definition. For the glb, take $\text{glb}(A) = \text{lub}\{x \in E \mid x \leq A\}$. This is Proposition 1.2 in Dowek's book. ■

Note that, if they exist, $\text{lub}\emptyset = \text{glb}E$ is the smallest element of E and $\text{lub}E = \text{glb}\emptyset$ the greatest.

A more abstract notion, not using sequences, is the one of *directed-complete partial order* (dCPO). This is the basis of *domain theory* used in programming languages semantics. We refer to [1] for more details. Domain theory has been introduced by Scott [5, 6, 8, 7].

Definition 4 (Directed-complete poset (dCPO)) A *directed set* D is a non-empty set such that every two elements has an upper bound (in D). A poset is *directed-complete* (dCPO) if every directed subset has a lub. In a dCPO, a function $f : E \rightarrow E$ is d-continuous (or Scott-continuous) if f is monotone and, for every directed subset D , $\text{lub}f(D) = f(\text{lub}D)$.

Note that, in the definition of d-continuity, $\text{lub}f(D)$ exists since, by monotony of f , $f(D)$ is directed.

By an easy induction, one can check that a directed set is a poset having an upper bound for every *finite* subset (including the empty subset since it is non-empty).

A chain is a directed set. Hence, every dCPO is a chain-complete poset (*i.e.* every chain has a lub) and thus an ω CPO. Conversely, using the axiom of choice, every chain-complete poset is a non-empty (or *pointed*) dCPO [4].

In Dowek's book, it is proved (Proposition 1.1, "1st fixed-point theorem") that, in an ω CPO, every ω -continuous function $f : E \rightarrow E$ has a least fixpoint (lfp), *i.e.* there is x such that $f(x) = x$. A similar theorem holds in dCPOs:

Lemma 5 In a dCPO, every d-continuous function $f : E \rightarrow E$ has a lfp.

In Dowek's book, it is also proved (Proposition 1.3, "2nd fixed-point theorem") that, in a complete lattice, every monotone function $f : E \rightarrow E$ has a lfp. This theorem is due to Tarski [10]. The full Tarski's theorem in fact says that the set of all the fixpoints of f is itself a complete lattice. And it can be extended to chain-complete posets [3].

Lemma 6 In a chain-complete poset, the set of fixpoints of a monotone function $f : E \rightarrow E$ is chain-complete. In particular, f has a lfp.

In fact, this property of monotone functions is a characteristic of chain-complete posets: if every monotone map $f : E \rightarrow E$ has a least fixpoint, then E is chain-complete [3].

2 Topology

In Dowek's book and in the previous section, the notion of limit of a chain or directed set is defined as its lub, and a monotone function $f : E \rightarrow E$ is continuous if it permutes with lub or if, for all directed subset D , $f(\text{lub}D) \leq \text{lub}f(D)$ (since we always have $\text{lub}f(D) \leq f(\text{lub}D)$ by monotony of f).

However, in mathematics, there exists a more abstract notion of continuity based on the notion of *topology*. Let $\wp(E)$ be the set of all the subsets of E .

Definition 7 (Topology) A *topology* τ on a set E is a set of subsets of E ($\tau \subseteq \wp(E)$), called *opens*, such that:

- \emptyset and E are opens: $\emptyset \in \tau$ and $E \in \tau$;
- τ is closed under arbitrary unions: if $X \subseteq \tau$, then $\bigcup X \in \tau$;
- τ is closed under *non-empty finite* intersections¹: if $X \subseteq \tau$ and X is *finite*, then $\bigcap X \in \tau$.

A function $f : E \rightarrow E$ is τ -*continuous* if the inverse image of an open is an open: for all $U \in \tau$, $f^{-1}(U) \in \tau$.

Alternative formulation using families:

- τ is closed under arbitrary unions: if $(U_i)_{i \in I}$ is an arbitrary family of opens, then $\bigcup_{i \in I} U_i$ is an open;
- τ is closed under *non-empty finite* intersections: if $(U_i)_{1 \leq i \leq n}$ is a *non-empty finite* family of opens (*i.e.* $n \geq 1$), then $\bigcap_{i \leq n} U_i$ is an open.

For instance, the standard notion of continuity on the real line \mathbb{R} corresponds to the following topology: $U \subseteq \mathbb{R}$ is open if, for all $x \in U$, there is $\epsilon > 0$ such that $]x - \epsilon, x + \epsilon[\subseteq U$. Hence, $]0, 1[$ is open, but $[0, 1]$ is not. Closure by *finite* intersections only is important: for all $i \geq 1$, $] - \frac{1}{i}, +\frac{1}{i}[$ is open, but

$\bigcap_{i \geq 1}] - \frac{1}{i}, +\frac{1}{i}[= \{0\}$ is not open.

We will see that the notion of d-continuity corresponds to the following topology:

¹Since τ has a greatest element E , one can define $\bigcap \emptyset$ as E and say that τ is closed under *finite* intersection (empty or not).

Definition 8 (Scott topology) Let σ be the set of subsets U such that:

- U is upper-closed: if $x \in U$ and $x \leq y$, then $y \in U$;
- U is accessible by directed lubs: for all directed subset D with $\text{lub}D \in U$, we have $D \cap U \neq \emptyset$.

We first check that σ is indeed a topology:

Lemma 9 σ is a topology.

Proof.

- One can easily check that \emptyset and E are upper-closed and accessible by directed lubs.
- Let (U_i) be a family of opens. Then, one can easily check that $\bigcup_{i \in I} U_i$ is upper-closed and accessible by directed lubs since each U_i do so.
- Let $(U_i)_{1 \leq i \leq n}$ be a non-empty finite family of opens. Then, one can easily check that $\bigcap_{1 \leq i \leq n} U_i$ is upper-closed. Let now D be a directed set such that $\text{lub}D \in \bigcap_{1 \leq i \leq n} U_i$. Then, for every $i \in \{1, \dots, n\}$, there is $x_i \in D \cap U_i$. Since D is directed, $x = \text{lub}\{x_1, \dots, x_n\} \in D$. Since every U_i is upper-closed, we have $x \in \bigcap_{1 \leq i \leq n} U_i$. ■

We now prove that d-continuity is equivalent to σ -continuity:

Lemma 10 A function $f : E \rightarrow E$ is d-continuous iff it is σ -continuous.

Proof. We first prove that d-continuity implies σ -continuity. Let $U \in \sigma$. We have to prove that $f^{-1}(U) \in \sigma$.

- $f^{-1}(U)$ is upper-closed: if $x \in f^{-1}(U)$ and $x \leq y$ then $f(x) \in U$ and, by monotony of f , $f(y) \in U$, i.e. $y \in f^{-1}(U)$.
- $f^{-1}(U)$ is accessible by directed lubs: let D be a directed subset such that $\text{lub}D \in f^{-1}(U)$. Then, $f(\text{lub}D) \in U$. By d-continuity, $f(\text{lub}D) = \text{lub}f(D)$. Since U is open, $f(D) \cap U \neq \emptyset$. Therefore, $D \cap f^{-1}(U) \neq \emptyset$.

We now prove that σ -continuity implies d-continuity. To this end, first remark that $]a, +\infty[= \{x \in E \mid x \not\leq a\}$ is an open:

- $]a, +\infty[$ is upper-closed: if $x \not\leq a$ and $x \leq y$, then $y \not\leq a$. Otherwise, by transitivity, $x \leq a$.
- $]a, +\infty[$ is accessible by directed lubs: if D is directed and $\text{lub}D \not\leq a$, then there is $d \in D$ such that $d \not\leq a$. Otherwise, $D \leq a$ and $\text{lub}D \leq a$, which is not possible.

We now prove that f is d-continuous:

- f is monotone: let x and y such that $x \leq y$. We have to prove that $f(x) \leq f(y)$. If $f(x) \not\leq f(y)$, then $f(x) \in U =]f(y), +\infty[$ and $x \in f^{-1}(U)$. Since f is d-continuous and U is open, $f^{-1}(U)$ is open and, in particular, upper-closed. Thus, $y \in f^{-1}(U)$ and $f(y) \not\leq f(y)$, which is not possible.
- For all directed subset D , $f(\text{lub}D) \leq \text{lub}f(D)$: if $f(\text{lub}D) \not\leq \text{lub}f(D)$, then $f(\text{lub}D) \in U =]\text{lub}f(D), +\infty[$ and $\text{lub}D \in f^{-1}(U)$. Since f is d-continuous and U is open, $f^{-1}(U)$ is open and, in particular, accessible by directed lubs. Thus, there is $d \in D \cap f^{-1}(U)$. Hence, $f(d) \in f(D) \cap U$ and $f(d) \not\leq \text{lub}f(D)$, which is not possible. ■

A subset F of E is *closed* if its complement in E , $F - E$, is open. The set of closed sets has the following closure properties, dual to those of open sets:

- \emptyset and E are closed sets²;
- the set of closed sets is closed under *finite* unions: if X is a *finite* set of closed sets, then $\bigcup X$ is a closed set;
- the set of closed sets is closed under arbitrary intersections: if X is a set of closed sets, then $\bigcap X$ is a closed set.

Alternatively, using families:

- the set of closed sets is closed under *finite* unions: if $(F_i)_{1 \leq i \leq n}$ is a finite family of closed sets, then $\bigcup_{1 \leq i \leq n} F_i$ is a closed set;
- the set of closed sets is closed under arbitrary intersections: if $(F_i)_{i \in I}$ is an arbitrary family of closed sets, then $\bigcap_{i \in I} F_i$ is a closed set.

Finally, note that a function f is τ -continuous if, for all closed set F , $f^{-1}(F)$ is closed.

References

- [1] S. Abramsky and A. Jung. Domain theory. In S. Abramsky, D. M. Gabbay, and T. S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, volume 3, pages 1–168. Clarendon Press, 1994. Corrected and expanded version available on A. Jung’s web page.
- [2] G. Dowek. *Proofs and Algorithms. An introduction to Logic and Computability*. Undergraduate Topics in Computer Science. Springer, 2011.
- [3] G. Markowsky. Chain-complete p.o. sets and directed sets with applications. *Algebra Universalis*, 6:53–68, 1976.

²Sets that are both open and closed are sometimes called “clopen”: \emptyset and E are clopen.

- [4] J. Mayer-Kalkschmidt and E. F. Steiner. Some theorems in set theory and applications in the ideal theory of partially ordered sets. *Duke Math. J.*, 31:287–289, 1964.
- [5] D. S. Scott. A type theoretic alternative to ISWIM, CUCH, OWHY. Manuscript. Published in [9], 1969.
- [6] D. S. Scott. Continuous lattices. In E. Lawvere, editor, *Toposes, Algebraic Geometry and Logic*, number 274 in Lecture Notes in Mathematics, pages 97–136. Springer, 1972.
- [7] D. S. Scott. Domains for denotational semantics. In *Proceedings of the 9th International Colloquium on Automata, Languages and Programming*, Lecture Notes in Computer Science 140, 1982.
- [8] D. S. Scott. Some ordered sets in computer science. In I. Rival, editor, *Ordered sets*, pages 677–718. D. Reidel, 1982.
- [9] D. S. Scott. A type theoretic alternative to ISWIM, CUCH, OWHY. *Theoretical Computer Science*, 121(411-440), 1993. Reprint of [5].
- [10] A. Tarski. A lattice-theoretical fixpoint theorem and its application. *Pacific Journal of Mathematics*, 5:285–309, 1955.